# Ruddy duck

Scan for more information



#### **Species Description**

Scientific name: Oxyura jamaicensis

Native to: North America and western South

America

Habitat: Lowland wetlands with lush emergent

vegetation and areas of open water

A small, compact duck with broad, short wings and narrow, stiff tail erect. The males are easily recognised by their distinctive blue bill, often chestnut held body and white face. The females are a dull brown colour, with a cream coloured face intersected by a dark horizontal line.

Introduced into wildfowl collections in the UK in the 1950s and natural dispersal into Northern Ireland has led to an estimated 25 - 30 pairs with a wintering population of around seventy to seventy five birds. Due to a going eradication programme their numbers may be significantly lower.


The ruddy duck is a serious threat to populations of white-headed duck (classified as vulnerable by the International Union for Conservation of Nature) in Europe through hybridisation. Eradication is being carried out in the UK to ensure the continued survival of the white-headed duck in Europe.


Under the Invasive Alien Species (Enforcement and Permitting) Order (Northern Ireland) 2019 it is offence to intentionally keep; breed; transport to, from or within Northern Ireland, use or exchange Ruddy duck; or to release it into the environment.



© Murray Foubister - CC BY-SA 2.0 - https://commons.wikimedia.org/w/index.php?curid=80621120

#### **Key ID Features**





Report any sightings via; CEDaR Online Recording - <a href="https://www2.habitas.org.uk/records/ISI">https://www2.habitas.org.uk/records/ISI</a>, iRecord app or Invasive Species Northern Ireland website - <a href="http://invasivespeciesni.co.uk/report-sighting">http://invasivespeciesni.co.uk/report-sighting</a>

#### Identification throughout the year

Ruddy duck are resident throughout the year in the UK. The plumage of females remains the same throughout the year. Between autumn and early spring the male's bill turns grey and its body colour becomes dull brown, however its white face remains. During the breeding season male ruddy duck are likely to be in the vicinity of any females and can be used to help identification, however, outside the breeding season individuals can occur on their own.


## Field Signs

The following field signs are shared by many species of duck and should not be used for the purpose of specific identification as ruddy duck.

- Call Generally silent.
- Nest In thick vegetation on water, creating a platform of reed or rush stems and leaves.
- Eggs 6-10 in clutch, dull or creamy white. Breeding period mid-April to July.


#### Distribution

Populations in GB have led to natural dispersal to Northern Ireland



### Similar Species

In the UK, male ruddy duck are unlikely to be confused with any other species. However, female ruddy duck are similar to females of a number of other duck species. The majority of female ducks are generally brown and are distinguished by a combination of size, body shape and subtle plumage characteristics.



Large bulbous bill

with some yellow

Mullarney, K, Svensson, L, Zetterstrom, D and Grant, PJ (2001) "Bird Guide" HarperCollins Snow, DW and Perrins, CM (1998) "The Birds of the Western Palearctic". Oxford University Press

Photos from: Len Blumin, Robin Ward, Lynn Watson.

**Sustainability** at the heart of a living, working, active landscape valued by everyone.



Brown body

44 – 45 cm

Unlikely to be

confused with

ruddy duck as usually found on

coastal waters



Adapted for Northern Ireland Environment Agency 2020

